

## **POST-OPERATIVE INSTRUCTIONS FOR SHOULDER ARTHROSCOPY**

(Acromioplasty and Debridement)

**PAIN CONTROL:** Take pain medication as directed. If you are experiencing a lot of pain at physical therapy, arrange your medication schedule to correspond to ½ hour before your scheduled appointment.

**ACTIVITY:** Rest the day of surgery. You may place a pillow under your forearm for comfort. It may be easier to sleep sitting up for the first few days.

**SLING:** The sling is only to help you feel more comfortable. You may take it off at any time.

**CRYOCUFF:** The cryocuff or ice wrap will be placed on your shoulder in the recovery room. Leave it on as much as possible for the first 3 days. After 3 days, use it for a minimum of 4 times a day for 30 minutes. The more you use it, the less swelling and inflammation you will have.

**BANDAGES:** You may remove the outer dressing and gauze pads after 48 hours. Do not remove the steri-strips (small pieces of tape running across the incision). You may cover the sutures loosely with a Band-Aid. Keep your sutures clean and dry. If you wish to shower, use a plastic bag or saran wrap to keep your shoulder dry. Your sutures will be removed approximately 10-14 days after surgery. Please call the office to schedule this appointment.

**PHYSICAL THERAPY:** Please contact the office to arrange physical therapy start-date.

**DRIVING:** No driving is allowed while you are wearing the sling.

**RETURNING TO WORK OR SCHOOL:** You may return to work (sedentary) or school 3-5 days after surgery if pain is tolerable. Returning to heavy labor will be determined by Dr. Allen.

---

### **NORMAL SENSATIONS AND FINDINGS AFTER SURGERY:**

- a. Shoulder pain
- b. Shoulder warmth up to 3 weeks
- c. Small amount of bloody drainage
- d. Numbness to incision area
- e. Bruising
- f. Swelling of the hand. If this happens, open and close your fingers to help pump the blood.
- g. Low-grade temperature less than 101.5 degrees – if this occurs, drink plenty of fluids and cough and deep breathe (take 10 breaths, on the last hold for a second then forcefully cough a few times). A low-grade temperature is normal for a week after surgery.
- h. Small amount of redness to the area where the sutures insert the skin

---

### **NOTIFY OUR OFFICE IMMEDIATELY IF ANY OF THE FOLLOWING SIGNS OR SYMPTOMS OCCUR:**

- a. Change is noted to your incision (i.e. increased redness or drainage)
- b. Temperature greater than 101.5 degrees
- c. Fever, chills, nausea, vomiting, or diarrhea
- d. Sutures become loose or fall out and incision becomes open
- e. Drainage becomes yellow, pus-like, or foul-smelling
- f. Increased pain unrelieved by medication or measures mentioned above